



Edinburgh Research Explorer

Ubiquitination of the human immunodeficiency virus type 1 Env glycoprotein

Citation for published version:

Bultmann, A, Eberle, J & Haas, J 2000, 'Ubiquitination of the human immunodeficiency virus type 1 Env glycoprotein', *Journal of Virology*, vol. 74, no. 11, pp. 5373-5376. <https://doi.org/10.1128/JVI.79.24.15074-15083.2005>

Digital Object Identifier (DOI):

[10.1128/JVI.79.24.15074-15083.2005](https://doi.org/10.1128/JVI.79.24.15074-15083.2005)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Virology

Publisher Rights Statement:

Copyright © 2000, American Society for Microbiology. All Rights Reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Ubiquitination of the Human Immunodeficiency Virus Type 1 Env Glycoprotein

ANDREAS BÜLTMANN, JOSEF EBERLE, AND JÜRGEN HAAS*

Max-von-Pettenkofer Institut, Genzentrum, LMU München, 81377 Munich, Germany

Received 12 October 1999/Accepted 7 March 2000

Expression of the human immunodeficiency virus type 1 (HIV-1) Env glycoprotein is stringently regulated in infected cells. The majority of the glycoprotein does not reach the cell surface but rather is retained in the endoplasmic reticulum or a *cis*-Golgi compartment and subsequently degraded. We here report that Env of various HIV-1 isolates is ubiquitinated at the extracellular domain of gp41 and that Env expression could be increased by lactacystin, a specific proteasome inhibitor, suggesting that the ubiquitin/proteasome system is involved in control of expression and degradation.

The human immunodeficiency virus type 1 (HIV-1) envelope glycoprotein gp160 is produced as a precursor polyprotein, which is cleaved in a Golgi or post-Golgi compartment by a cellular furin-type protease into gp120 and gp41 subunits (13, 19, 22, 25, 26). Although Env cleavage efficacy during its transport through the secretory pathway depends on both cell type and virus isolate, it is very inefficient; the majority of the Env glycoproteins remain uncleaved and retained in the endoplasmic reticulum (ER) or a *cis*-Golgi compartment (8). Accordingly, the majority of the gp160 glycoproteins remain endoglycosidase H sensitive, which is indicative of proteins that do not reach the medial Golgi (14, 27). Retained and probably misfolded Env is subsequently degraded, which was reported to occur in either lysosomes (31) or a Golgi-associated compartment (14). However, the current view of protein degradation holds that lysosomes are reached through either endosomes or late Golgi compartments.

The ER is the entry site of proteins into the secretory pathway. It is responsible for proper folding of proteins before transport to the *cis*-Golgi compartment and acts to ensure that misfolded and nonassembled proteins are eliminated (reviewed in references 10 and 15). Recently, the ubiquitin/proteasome system located in the cytoplasm was identified as the major site of degradation for ER-resident proteins as well as proteolytic substrates of the secretory pathway destined for degradation (3, 28). Proteolytic degradation of these proteins appears to be preceded by a retrograde transport to the cytosol (17, 30). Recently, it was reported that the chaperone GRB78 BiP is linked to this retrograde protein translocation for ER degradation (12, 18, 23). In HIV-infected cells, GRB78 BiP was shown to bind to Env (9) together with other ER-based chaperones including calnexin (16, 21) and calreticulin (21). As the majority of Env is retained in the ER and subsequently degraded without reaching late Golgi compartments, we asked whether the ubiquitin/proteasome system is involved in HIV-1 Env degradation and whether ubiquitinated Env can be detected in infected cells.

First, we tested whether Env glycoprotein of the MN isolate can be reprecipitated using antiubiquitin and anti-Env antibodies (Fig. 1a). Upon sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) Env glycoprotein was detected




FIG. 1. Env from various HIV-1 subtypes is ubiquitinated. (a) 293T cells transiently transfected with Env MN were pulsed for 1 h with [³⁵S]Cys-Met, lysed, and precipitated with either polyclonal antiserum U5379 against ubiquitin (Sigma, St. Louis, Mo.), a human antiserum detecting HIV-1 Env (95-2), or an irrelevant antiserum. Subsequently, lysates were reprecipitated with either of the three antibodies and analysed by SDS-PAGE (8% gel) and autoradiography. The numbers 1 and 2 refer to the sequential order of precipitation. (b) Ubiquitinated Env of cells metabolically labeled for 1 h with [³⁵S]Cys-Met can be detected by precipitation using monoclonal antibody 2C5 (Calbiochem, La Jolla, Calif.) or polyclonal antisera SPA-200 (Stressgene, Victoria, British Columbia, Canada) and U5379 (Sigma) against ubiquitin and reprecipitation with anti-Env. (c) 293T cells were either infected with recombinant vaccinia virus (NIBSC/MRC AIDS Reagent Project) expressing Env of the 92UG037 (subtype A), 92BR020 (subtype B), or 92BR025 (subtype C) isolates or transiently transfected with a plasmid expressing Env of the MN (subtype B) isolate (1). Cell lysates of cells metabolically labeled for 1 h with [³⁵S]Cys-Met were either precipitated with human antiserum 95-2 against HIV-1 or first precipitated with polyclonal serum U5379 against ubiquitin and reprecipitated with antiserum 95-2 (derived from an HIV-1-infected individual). Immunoprecipitates were subjected to SDS-PAGE on an 8% gel. (d) Cells transiently transfected with Env MN were labeled for 1 h with [³⁵S]Cys-Met, lysed, and precipitated with either antiserum 95-2 or polyclonal antiserum U5379 and subsequently reprecipitated with 95-2. The immunoprecipitates were either digested with PNGase F (Roche, Mannheim, Germany) (gp160*) overnight or left untreated (gp160) and subjected to electrophoresis.

* Corresponding author. Mailing address: Max-von-Pettenkofer Institut, Genzentrum, LMU München, Feodor-Lynen-Str. 25, 81377 Munich, Germany. Phone: 49 89 2180 6852. Fax: 49 89 2180 6899. E-mail: haas@lmb.uni-muenchen.de.




FIG. 2. Ubiquitinated Env is membrane associated, and its expression is increased by proteasome inhibitors. Transiently transfected 293T cells expressing Env MN were pulsed for 1 h with [35 S]Cys-Met and subsequently chased for 0 and 5 h. Cells were pretreated with 5 μ M lactacystin (+) or culture medium alone (-) for 1 h. Lactacystin was also present during the pulse and chase periods. Cells were detached with 1 mM EDTA-phosphate-buffered saline (PBS) and resuspended in hypotonic buffer (10 mM HEPES, 10 mM KCl, 10 mM MgCl₂ [pH 7.6]). Subsequently, cells were disrupted using a 25-gauge syringe, nuclei were removed by centrifugation at $1,000 \times g$ for 10 min, and membrane (m) and cytosolic (c) fractions were isolated by centrifugation at $20,000 \times g$ for 15 min. The pellet was resuspended and washed once in hypotonic buffer. The lysates were split and either precipitated with human antiserum 95-2 against HIV-1 (a), first precipitated with a polyclonal serum against ubiquitin (U5379) and reprecipitated with antiserum 95-2 (b), or precipitated with a monoclonal antibody against MHC class I (W6/32) (c). Immunoprecipitates were subjected to electrophoresis on either an 8% (gp160) or 12% (MHC class I) gel.

independently of the sequential order of precipitation if anti-ubiquitin and anti-Env antibodies were used, but not if either of the two antibodies was replaced by an irrelevant antibody. To exclude that cross-reactivity of the antiubiquitin antibody caused this result, we tested three different commercially available mono- and polyclonal antibodies (Fig. 1b). As shown with Env 92BR020, all three antibodies were able to reprecipitate Env, indicating specific recognition of the antigen. Subsequently, we tested glycoproteins of different HIV-1 isolates expressed by either recombinant vaccinia virus or transient transfection with Env expression plasmids for ubiquitination (Fig. 1c). Env glycoproteins from all HIV-1 isolates tested, including MN (subtype B) and the primary isolates 92UG037 (subtype A), 92BR020 (subtype B), and 92BR025 (subtype C), were found to be ubiquitinated. Moreover, ubiquitinated Env was precipitated from 293T kidney carcinoma cells (Fig. 1), as well as C8166 and Jurkat T lymphocytes, indicating that ubiquitination of Env is not cell type specific (data not shown). Although the molecular weight was difficult to evaluate due to high glycosylation, we could detect no major differences in size between total Env and ubiquitinated Env, suggesting that only a single ubiquitin molecule (8.4 kDa) had been added. To evaluate the number of ubiquitin residues linked to the glycoprotein, we treated the immunoprecipitates with peptide *N*-glycosidase (PNGase F), which completely removes carbohydrate moieties (Fig. 1d). A size difference of approximately 8 kDa corresponding to one ubiquitin molecule could be detected between deglycosylated total and ubiquitinated Env. No ubiquitinated Env could be found within total deglycosylated Env, probably reflecting the low quantity of the ubiquitinated form in relation to total Env.

Misfolded soluble and membrane proteins in the ER are

dislocated to the cytosolic proteolytic system by retrograde transport, in which at least for some proteins components of the Sec61 translocon and the proteasome appear to be involved (17, 30). We thus tested by particulate fraction experiments whether ubiquitinated Env is membrane associated or can be found in the cytosol (Fig. 2). Transfected 293T cells were metabolically labeled for 1 h and chased for 0 and 5 h. Cells were disrupted and separated into membrane and cytosolic fractions by centrifugation. After both 0 and 5 h of chase, ubiquitinated Env was found only in the membrane (pellet), not in the cytosolic fraction (supernatant). Due to the low Env cleavage efficiency in 293T cells, almost no gp120 could be detected even after 5 h of chase. To control for proper fractionation, major histocompatibility complex (MHC) class I molecules were precipitated from both fractions. As previously reported, MHC class I molecules were precipitated from the membrane but not in the cytosolic fraction under the conditions used (30). Therefore, we have no evidence for a contamination of the two fractions and conclude that the main portion of ubiquitinated Env is still membrane bound and not located within the cytosol in soluble form. Addition of lactacystin, a specific proteasome inhibitor, increased the amount of both Env and ubiquitinated Env after 0 and 5 h of chase (Fig. 2). Thus, these data suggest that ubiquitinated Env is at least partly degraded by proteasomes and that Env is ubiquitinated prior to translocation into the cytosol.

Ubiquitin is usually attached to lysine residues in long polymeric chains (7, 11) but can also be detected in monoubiquitinated (20, 29) and in lysineless proteins (5, 6). To determine the site of ubiquitination, C-terminal Env deletion mutants were tested by reprecipitation with antiubiquitin and anti-Env antibodies (Fig. 3a and b). Ubiquitinated Env MN was detected in transfected 293T cells after a 1-h pulse if gp160, gp140 Δ cyt, or gp130 Δ tm was expressed, but not for gp120, suggesting that ubiquitin is bound to the extracellular part of gp41. Further evidence for this binding site was received by proteinase K digest of isolated microsomes from 293T cells expressing Env (Fig. 3c). Ubiquitinated Env was detected by reprecipitation in undigested as well as in proteinase K-digested microsomes, indicating that proteolytic deletion of the cytoplasmic Env domain does not disrupt ubiquitination. Since we were not able to detect the proteinase K-digested Env protein by Western blot analysis with monoclonal antibody 1577 directed against residues 735 to 752 of gp41, the cytoplasmic gp41 domain had been completely degraded (Fig. 3d). Sequence comparison between the four ubiquitinated Env glycoproteins shown in Fig. 1 revealed three conserved lysine residues as potential sites of ubiquitination. Our results indicating that ubiquitinated Env is still membrane associated and that ubiquitin is bound to the extracellular, luminal part of gp41 suggest that ubiquitination, translocation, and ER degradation of Env constitute an integrated process, as recently discussed for other luminal and integral membrane proteins (4, 24). To test whether ubiquitinated Env can also be detected in HIV-1-infected cells, we metabolically labeled C8166 cells infected with either HIV-1 MN or HIV-1 MvP 899 and reprecipitated cell lysates with antiubiquitin and anti-Env antisera (Fig. 4). In accordance with the experiments shown in Fig. 3, mapping the ubiquitination site, we detected ubiquitinated gp160 but not gp120 in cells infected with both HIV-1 isolates.

In this study, we showed that the extracellular domain of HIV-1 gp41 is ubiquitinated. Since the proteasome inhibitor lactacystin increased the amount of Env, we assume that ubiquitination occurs prior to degradation in the proteasome. This is consistent with and might explain observations by us and other groups that the majority of Env molecules are retained in




FIG. 3. The extracellular domain of gp41 is ubiquitinated. (a) Schematic diagram of the Env MN deletion mutants used. (b) Env MN deletion mutants were transiently transfected into 293T cells. Cell lysates of metabolically labeled cells were either precipitated with human antiserum 95-2 against Env or first precipitated with polyclonal serum U5379 (Sigma) against ubiquitin and reprecipitated with antiserum 95-2. Immunoprecipitates were subjected to SDS-PAGE on an 8% gel. (c) 293T cells transiently transfected with Env MN were metabolically labeled for 1 h. Subsequently, cells were detached with 1 mM EDTA-PBS, washed with PBS, resuspended in lysis buffer (140 mM NaCl, 5 mM MgCl₂, 20 mM Tris-HCl [pH 7.6]) without NP-40, and homogenized in a cell homogenizer. Nondisrupted cells and nuclei were removed by low-speed centrifugation, and microsomes were isolated by centrifugation for 1 h at 100,000 × g. The microsome fraction was resolved and subjected to either proteinase K (0.4 mg/ml, final concentration) or proteinase K and NP-40 (1%) digestion for 30 min. Full-length Env (*) or Env lacking the cytoplasmic domain (**) was either precipitated with human antiserum 95-2 against HIV-1 or first precipitated with a monoclonal antibody against ubiquitin and reprecipitated with antiserum 95-2. Immunoprecipitates were subjected to SDS-PAGE on an 8% gel. (d) Untreated, proteinase K-treated, or proteinase K- and NP40-treated microsomal fractions were analyzed by Western blotting using human antiserum 95-2 against Env or monoclonal antibody 1577 directed against the cytoplasmic domain of gp41.

the ER and stay endoglycosidase H sensitive (14, 27). Intriguingly, recent reports suggested alternative explanations for ubiquitinated viral proteins. It was shown that p6^{gag} of HIV-1 and simian immunodeficiency virus, as well as p12^{gag} of murine

leukemia virus and US9 of herpes simplex virus type 1 are ubiquitinated and can be found within virus particles, in addition to free ubiquitin (2, 5, 20). The significance of ubiquitinated viral proteins thus remains unclear. Ubiquitinated viral proteins might have escaped from degradation by the proteasome in cells with high levels of viral proteins. Alternatively, it has been suggested that ubiquitinated viral proteins have specific roles in viral pathogenesis using the host cell ubiquitin/proteasome machinery (5, 20).

This work was supported by the Sonderforschungsbereich 464 (Teilprojekt B6) of the Deutsche Forschungsgemeinschaft.

Vaccinia virus recombinants were provided by M. Esteban, and monoclonal antibody 1577 was provided by M. Ferguson through courtesy of the NIBSC/MRC AIDS Reagent Project. We thank H. Ziegler for advice and reading of the manuscript.

REFERENCES

1. Andre, S., B. Seed, J. Eberle, W. Schraut, A. Bultmann, and J. Haas. 1998. Increased immune response elicited by DNA vaccination with a synthetic gp120 sequence with optimized codon usage. *J. Virol.* **72**:1497-1503.
2. Arthur, L. O., J. W. Bess, Jr., R. C. Sowder, R. E. Benveniste, D. L. Mann, J. C. Chermann, and L. E. Henderson. 1992. Cellular proteins bound to immunodeficiency viruses: implications for pathogenesis and vaccines. *Science* **258**:1935-1938.
3. Biederer, T., C. Volkwein, and T. Sommer. 1996. Degradation of subunits of the Sec61p complex, an integral component of the ER membrane, by the ubiquitin-proteasome pathway. *EMBO J.* **15**:2069-2076.
4. Biederer, T., C. Volkwein, and T. Sommer. 1997. Role of Cuelp in ubiquitination and degradation at the ER surface. *Science* **278**:1806-1809.




FIG. 4. gp160 is ubiquitinated in HIV-1-infected cells. C8166 cells infected with either HIV-1 MN or HIV-1 MvP 899 or mock-infected control cells were metabolically labeled with [³⁵S]Cys-Met for 15 h and subsequently lysed without chase. Cell lysates were either precipitated with human antiserum 95-2 against Env or first precipitated with polyclonal serum U5379 (Sigma) against ubiquitin and reprecipitated with antiserum 95-2.

5. Brandimarti, R., and B. Roizman. 1997. Us9, a stable lysine-less herpes simplex virus 1 protein, is ubiquitinated before packaging into virions and associates with proteasomes. *Proc. Natl. Acad. Sci. USA* **94**:13973–13978.
6. Breitschopf, K., E. Bengal, T. Ziv, A. Admon, and A. Ciechanover. 1998. A novel site for ubiquitination: the N-terminal residue, and not internal lysines of MyoD, is essential for conjugation and degradation of the protein. *EMBO J.* **17**:5964–5973.
7. Ciechanover, A. 1994. The ubiquitin-proteasome proteolytic pathway. *Cell* **79**:13–21.
8. Crise, B., L. Buonocore, and J. K. Rose. 1990. CD4 is retained in the endoplasmic reticulum by the human immunodeficiency virus type 1 glycoprotein precursor. *J. Virol.* **64**:5585–5593.
9. Earl, P. L., B. Moss, and R. W. Doms. 1991. Folding, interaction with GRP78-BiP, assembly, and transport of the human immunodeficiency virus type 1 envelope protein. *J. Virol.* **65**:2047–2055.
10. Hammond, C., and A. Helenius. 1995. Quality control in the secretory pathway. *Curr. Opin. Cell Biol.* **7**:523–529.
11. Jentsch, S., and S. Schlenker. 1995. Selective protein degradation: a journey's end within the proteasome. *Cell* **82**:881–884.
12. Joslin, G., W. Hafeez, and D. H. Perlmuter. 1991. Expression of stress proteins in human mononuclear phagocytes. *J. Immunol.* **147**:1614–1620.
13. Kantanen, M. L., P. Leinikki, and E. Kuusmanen. 1995. Endoproteolytic cleavage of HIV-1 gp160 envelope precursor occurs after exit from the trans-Golgi network (TGN). *Arch. Virol.* **140**:1441–1449.
14. Kimura, T., M. Nishikawa, and J. Fujisawa. 1996. Uncleaved env gp160 of human immunodeficiency virus type 1 is degraded within the Golgi apparatus but not lysosomes in COS-1 cells. *FEBS Lett.* **390**:15–20.
15. Kopito, R. R. 1997. ER quality control: the cytoplasmic connection. *Cell* **88**:427–430.
16. Li, Y., J. J. Bergeron, L. Luo, W. J. Ou, D. Y. Thomas, and C. Y. Kang. 1996. Effects of inefficient cleavage of the signal sequence of HIV-1 gp 120 on its association with calnexin, folding, and intracellular transport. *Proc. Natl. Acad. Sci. USA* **93**:9606–9611.
17. Mayer, T. U., T. Braun, and S. Jentsch. 1998. Role of the proteasome in membrane extraction of a short-lived ER-transmembrane protein. *EMBO J.* **17**:3251–3257.
18. Meerovitch, K., S. Wing, and D. Goltzman. 1998. Parathyroid hormone-related protein is associated with the chaperone protein BiP and undergoes proteasome-mediated degradation. *J. Biol. Chem.* **273**:21025–21030.
19. Merkle, R. K., D. E. Helland, J. L. Welles, A. Shilatfard, W. A. Haseltine, and R. D. Cummings. 1991. gp160 of HIV-1 synthesized by persistently infected Molt-3 cells is terminally glycosylated: evidence that cleavage of gp160 occurs subsequent to oligosaccharide processing. *Arch. Biochem. Biophys.* **290**:248–257.
20. Ott, D. E., L. V. Coren, T. D. Copeland, B. P. Kane, D. G. Johnson, R. C. Sowder, Y. Yoshinaka, S. Oroszlan, L. O. Arthur, and L. E. Henderson. 1998. Ubiquitin is covalently attached to the p6^{Gag} proteins of human immunodeficiency virus type 1 and simian immunodeficiency virus and to the p12^{Gag} protein of Moloney murine leukemia virus. *J. Virol.* **72**:2962–2968.
21. Otteken, A., and B. Moss. 1996. Calreticulin interacts with newly synthesized human immunodeficiency virus type 1 envelope glycoprotein, suggesting a chaperone function similar to that of calnexin. *J. Biol. Chem.* **271**:97–103.
22. Pfeiffer, T., H. Zentgraf, B. Freyaldenhoven, and V. Bosch. 1997. Transfer of endoplasmic reticulum and Golgi retention signals to human immunodeficiency virus type 1 gp160 inhibits intracellular transport and proteolytic processing of viral glycoprotein but does not influence the cellular site of virus particle budding. *J. Gen. Virol.* **78**:1745–1753.
23. Plemper, R. K., S. Bohmler, J. Bordallo, T. Sommer, and D. H. Wolf. 1997. Mutant analysis links the translocon and BiP to retrograde protein transport for ER degradation. *Nature* **388**:891–895.
24. Plemper, R. K., R. Egner, K. Kuchler, and D. H. Wolf. 1998. Endoplasmic reticulum degradation of a mutated ATP-binding cassette transporter Pdr5 proceeds in a concerted action of Sec61 and the proteasome. *J. Biol. Chem.* **273**:32848–32856.
25. Raja, N. U., M. J. Vincent, and M. A. Jabbar. 1993. Analysis of endoproteolytic cleavage and intracellular transport of human immunodeficiency virus type 1 envelope glycoproteins using mutant CD4 molecules bearing the transmembrane endoplasmic reticulum retention signal. *J. Gen. Virol.* **74**:2085–2097.
26. Salzwedel, K., J. T. West, Jr., M. J. Mulligan, and E. Hunter. 1998. Retention of the human immunodeficiency virus type 1 envelope glycoprotein in the endoplasmic reticulum does not redirect virus assembly from the plasma membrane. *J. Virol.* **72**:7523–7531.
27. Stein, B. S., and E. G. Engleman. 1990. Intracellular processing of the gp160 HIV-1 envelope precursor. Endoproteolytic cleavage occurs in a cis or medial compartment of the Golgi complex. *J. Biol. Chem.* **265**:2640–2649.
28. Ward, C. L., S. Omura, and R. R. Kopito. 1995. Degradation of CFTR by the ubiquitin-proteasome pathway. *Cell* **83**:121–127.
29. West, M. H., and W. M. Bonner. 1980. Histone 2B can be modified by the attachment of ubiquitin. *Nucleic Acids Res.* **8**:4671–4680.
30. Wiertz, E. J., D. Tortorella, M. Bogoy, J. Yu, W. Mothes, T. R. Jones, T. A. Rapoport, and H. L. Ploegh. 1996. Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature* **384**:432–438.
31. Willey, R. L., J. S. Bonifacino, B. J. Potts, M. A. Martin, and R. D. Klausner. 1988. Biosynthesis, cleavage, and degradation of the human immunodeficiency virus 1 envelope glycoprotein gp160. *Proc. Natl. Acad. Sci. USA* **85**:9580–9584.